





EASTERN REGION
STATE FOREST LANDS
ANNUAL WORK PLAN
FISCAL YEAR 2017

Prepared:		4-4-16
	(Forest Manager)	Date
Reviewed:		4-6-16
	(Regional Manager)	Date
Reviewed:		5/5/16
	(Land Acquisition & Planning)	Date
Approved:		5.09.16
	(Environmental Specialist)	Date

Prepared By:

Michael G. Schofield, MFS – Chesapeake Forest Manager

Alexander Clark, MFS – GIS Forester

Contributors:

Skip Jones, Parker Forestry Services Inc.

DNR Interdisciplinary Team

Citizens Advisory Committee



The mark of
responsible forestry

CONTENTS

A. Forest Overview	4
Chesapeake Forest and Pocomoke State Forest	4
Historic Forest Conditions and the Role of Fire	4
Forest Types and Size Classes	5
Unique Community Types	6
B. Annual Work Plan Summary	8
Introduction	8
Networking with DNR and other agencies	8
C. Maintenance Projects	9
D. Recreation Projects	9
E. Special Projects	10
F. Watershed Improvement Projects	10
G. Special Wildlife Habitat Projects	10
H. Ecosystem Resoration Projects	11
I. Monitoring Projects	11
J. Review Process	12
Interdisciplinary Team Comments	12
Citizen's Advisory Committee Comments	21
Public Comments	24
K. Silvicultural Projects	28
Silvicultural Activity Overview	28
Definitions of Silvicultural Activities	28
Silvicultural Prescriptions & Stand Data	31
Caroline County	31
Dorchester County	31
Somerset County	33
Wicomico County	34
Worcester County	37
Pocomoke State Forest	39
Silvicultural Site Maps	40
L. Budget	66
Appendix A - Recreation Trail Grants	67
Appendix B - Soil Series Management Groups, Abbreviations, and Symbols	86
Chesapeake Forest/Pocomoke State Forest: Soil Management Groups	88
Appendix C. Silvicultural Activity Summaries	90
Works Cited	93

A. FOREST OVERVIEW

CHESAPEAKE FOREST AND POCOMOKE STATE FOREST

The Chesapeake Forest which is owned by the State of Maryland and managed by the Maryland Forest Service through the Department of Natural Resources originally consisted of 58,000 acres of forest land. These lands were part of a 1999 divestment by the Chesapeake Forest Products Corporation. At that time, a partnership between the State of Maryland, The Conservation Fund, and Hancock Timber Resources Group moved to purchase the forests. The original 1999 plan was prepared by a 10-person technical team assembled by The Sampson Group, Inc. Oversight and decision making for the technical team was provided by a Steering Committee composed of representatives from Maryland Department of Natural Resources, The Conservation Fund, the Chesapeake Bay Foundation, and the local forest industry.

The Chesapeake Forest currently consists of 67,731 acres divided into 185 Management Units distributed across six counties. Chesapeake Forest also includes the Seth Demonstration Forest in Talbot County, Wicomico Demonstration Forest in Wicomico County, and Fred W. Besley Demonstration Forest in Dorchester County. In spite of this scattered character, the forests include some of the last large segments of unbroken forest in a region that is largely agricultural in nature. Chesapeake Forest Lands include more than 6,000 acres of wetlands or swamps and comprise portions of 23 separate watersheds, many of which have been given a high priority for conservation action under the Maryland Clean Water Action Plan. They contain established populations of threatened and endangered species, including the Delmarva fox squirrel (*Sciurus niger cinereus*), bald eagle, and some 150 other species that have been identified as rare, threatened, or endangered in the region. Abundant populations of deer, turkey, and waterfowl create the basis for extensive hunting opportunities and other recreational activities on the land.

The 17,918-acre Pocomoke State Forest is almost entirely contained within Worcester County, except for 429 acres in Somerset County and 145 acres in Wicomico County. The Chesapeake Forest has 17,613 acres within Worcester County, and several tracts from both Chesapeake Forest and Pocomoke State Forest adjoin each other offering greater habitat and recreational management opportunities. In addition, since both forests contain similar forest types, many of the same management guidelines and principles are used. There are differences between the two forests, however. Pocomoke State Forest contains many older tracts of forestland still in their natural state, nearly 5,000 acres of cypress and hardwood forest that borders a state scenic river, and areas of state designated Wildlands.

For additional information about Chesapeake Forest and Pocomoke State Forest please visit their respective web pages located at: <http://www.dnr.state.md.us/forests/mdforests.asp>.

HISTORIC FOREST CONDITIONS AND THE ROLE OF FIRE

The average pre-European-settlement fire frequency was on the order of 7-12 years for forests of the Eastern Shore of Maryland, with higher frequencies of 4-6 years in the southeastern Maryland counties of Wicomico, Worcester, Somerset, and Dorchester (Frost, 1998). These frequencies are high compared to most areas of the Northeast. Since it is unlikely that lightning was a significant contributor to these fires, Native American populations must have been. A conclusion is that fire in the Northeast was predominantly a phenomenon associated with human activity (Pyne, 1982).

The forest that covered the Eastern Shore in Indian times was primarily a hardwood one, though increasingly mixed with pine to the southward (Rountree & Davidson, 1997). The large patches of pine-dominated woods today are largely second growth, the result of extensive clearing in historic times. In aboriginal times, the woods of the Eastern Shore were likely to be oak-hickory, oak-gum, or oak-pine types, all of which still exist in second-growth form.

Captain John Smith said in the early seventeenth century, “A man may gallop a horse amongst these woods any waie, but where the creekes or Rivers shall hinder”. Father Andrew White wrote that the woods around St. Mary’s were so free of underbrush that a “coach and fower horses” could be driven through them (Rountree & Davidson, 1997). The open conditions could be partly attributed to the closed canopies of these mature forests, which shaded out undergrowth, but it is also likely that periodic fire helped to maintain the park-like conditions.

It is reasonable to assume that Eastern Shore tribes also used fire to periodically burn the marshes that were important sources of mollusks, fish, furbearers, waterfowl, edible tubers, and reeds for housing. Fire would have been useful for herding game, enhancing visibility or access, or retarding invasion of woody growth. More often than not, these fires would have spread into adjacent woodlands and, if of sufficient intensity, created the open seedbed conditions conducive to establishment of loblolly pine. Even today the pattern of loblolly pine “islands” and “stringers” in and adjacent to marshes of the lower Eastern Shore is common.

If, as Rountree and Davidson suggest, oaks were the most prevalent species in pre-settlement times, then the possible role of fire in maintaining these forest types must also be considered. Frost stated, “Light, understory fires may have been the norm for millions of hectares of eastern hardwood forest...” (Frost, 1998). Oak species range from slightly tolerant to intolerant of shade, indicating that disturbance is desirable to promote regeneration and growth. Furthermore, acorn germination and initial seedling establishment are most successful where light understory burns have scarified the seedbed and reduced competition (Burns & Honkala, 1990). The extensive presence of oaks on the Shore was an indicator that low-intensity understory fires were common, either intentionally set by Indians to create “open woods” or drive game, or the incidental result of land-clearing.

Natural stands of loblolly pine (*Pinus taeda*) became much more widespread around the turn of the 20th Century, particularly in the counties south of the Choptank River, largely due to the influence of economic factors. First was the abandonment of agricultural fields as farmers moved to more lucrative jobs in the towns and cities. Loblolly pine is an opportunistic species, which found the recently abandoned fields prime sites for reproduction by natural seeding. The second factor was the rise of large-scale commercial lumbering. Steam locomotives, often used to haul logs from the woods, were notorious for throwing sparks along the tracks and starting fires. Both the clearing of the forests by large-scale logging and the subsequent fires resulted in large areas of open, scarified land suitable for pine regeneration. By the middle of the twentieth century, loblolly pine had become the predominant forest cover type in the lower counties of the Eastern Shore.

FOREST TYPES AND SIZE CLASSES

Young loblolly pine forests mostly established since the early 1980’s are what characterize a high proportion of the Chesapeake Forest. Mixed pine and hardwood forests still occupy some of the lands, and many riparian areas and flood plains contain stands of mixed hardwoods. In general, the mixed pine-hardwood and hardwood stands are older, mature forests.

Mature mixed pine-hardwood, bottomland hardwood, and bald-cypress forests comprise the majority of the Pocomoke State Forest. In general, the mixed pine-hardwood, hardwood, and bald cypress stands are older, mature forests, while loblolly pine stands are more evenly distributed across all age classes.

Table 1 provides a habitat diversity matrix of both Eastern Region State Forests that provides a current baseline from which future changes in age structure or forest type diversity can be assessed for potential habitat or biodiversity effects.

Table 1. Forest Diversity Analysis

Acres of forest type and forest structure by structural groups, with percent of total area in each forest type/structure group combination.

Forest type	Structure stage							Total Area
	Open 0 - 5 yrs	Sapling 5 - 15 yrs	Growing 15 - 25 yrs	Maturing 25 - 35 yrs	Mature 35 - 50 yrs	Big Trees 50 - 75+ yrs	Uneven Aged	
Atlantic White Cedar	4	3	0	0	0	0	0	7
(Percent)	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%	0.01%
Loblolly Pine	1,185	9,557	21,016	12,644	7,312	1,617	407	53,737
(Percent)	1.40%	11.28%	24.81%	14.93%	8.63%	1.91%	0.48%	63.44%
Shortleaf Pine	0	0	0	0	0	255	0	255
(Percent)	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%	0.30%
Mixed Pine/ Hardwood	721	886	933	717	1,563	7,568	22	12,410
(Percent)	0.85%	1.05%	1.10%	0.85%	1.85%	8.94%	0.03%	14.65%
Mixed Hardwoods	439	296	237	101	200	9,188	12	10,471
(Percent)	0.52%	0.35%	0.28%	0.12%	0.24%	10.85%	0.01%	12.36%
Bottomland Hardwoods/ Bald Cypress	0	0	0	0	20	3,855	0	3,875
(Percent)	0.00%	0.00%	0.00%	0.00%	0.02%	4.55%	0.00%	4.57%
Marsh/Field/ Power lines	3,946	0	0	0	0	0	0	3,946
(Percent)	4.66%	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%	4.66%
Total	6,295	10,741	22,186	13,462	9,095	22,483	441	85,533
(Percent)	7.43%	12.68%	26.19%	15.89%	10.74%	26.54%	0.52%	100.00%

UNIQUE COMMUNITY TYPES

Xeric or inland sand dunes are found primarily in the lower Eastern Shore counties. They are located on very well drained sand ridges formed by winds blowing off receding glaciers. These sand ridges support a variety of rare and threatened insect and plant species. The primary species in this community are shortleaf pine (*Pinus echinata*), Virginia pine (*Pinus virginiana*), and various oak species (*Quercus spp.*), with an understory comprised of lowbush blueberry (*Vaccinium pallidum*) and an assortment of ericaceous plants. Xeric sand dunes have been identified and mapped either as an Ecologically Significant Area (ESA) or as a Globally Rare (G3) Community.

Pond pine (*Pinus serotina*) forests are typically found in swamps and other poorly drained areas. Pond pine can be found along with pitch and loblolly pine, and it can hybridize with those species. During periods of drought, these forests can be subject to intense fires. Pond pine needs fire to open the serotinous cones and release the seeds to facilitate natural regeneration.

Delmarva bays and associated life zones are isolated depressional wetlands that serve the needs of wetland breeding animals and support several species of rare plants. Delmarva bays can vary in their ecological quality,

primarily due to past management practices. The hydrology of many bays was altered for agriculture or to attempt to increase forest production. Therefore, many of these bays may require restoration to get the bay back to a more natural state. Delmarva bays and the associated life zone have their own ESA designations identified and mapped.

Riparian swamps

Atlantic white cedar (*Chamaecyparis thyoides*) swamps are nontidal forests that border on rivers or headwaters of streams.

Bald cypress (*Taxodium distichum*) swamps and forests can be tidal or nontidal. These forests are known for their pronounced microtopography of hollows and hummocks.

Vernal pools and seasonal wetlands are temporary wetlands present in late winter and spring that support amphibian reproduction. These can be found throughout the eastern shore region.

B. ANNUAL WORK PLAN SUMMARY

INTRODUCTION

This section summarizes the proposed activities that will occur on all public forest lands (84,762 acres) managed by the Maryland Forest Service within the Eastern Region during the 2017 fiscal year. These lands include the Chesapeake Forest, Pocomoke State Forest, Wicomico Demonstration Forest, Seth Demonstration Forest, and Fred W. Besley Demonstration Forest. The fiscal year runs from July 1, 2016 to June 30, 2017. The following proposed activities are the results of a multi-agency effort. The multi-agency approach has ensured that all aspects of these lands have been addressed within the development of this plan.

All projects and proposals within this Plan have been developed to meet one or more of the Land Management Guidelines and Objectives as seen in the Chesapeake Forest and Pocomoke State Forest Sustainable Forest Management Plans including:

- **Forest Economy** - management activities with a purpose to maintain an economically sustainable forest and contribute to the local economy through providing forest-related employment and products.
- **Forest Conservation** - management activities with a purpose to protect significant or unique natural communities and elements of biological diversity, including Ecologically Significant Areas, High Conservation Value Forests and old growth Forests. Old growth forest management serves to restore and/or enhance old growth forest structure and function.
- **Water Quality** - management activities designed to protect or improve ecological functions in protecting or enhancing water quality.
- **Wildlife Habitat** - management activities with a purpose to maintain and enhance the ecological needs of the diversity of wildlife species and habitat types.
- **Recreation and Cultural Heritage** - management activities with a purpose to maintain and enhance areas that serve as visual, public camping, designated trails, and other high public use areas.

NETWORKING WITH DNR AND OTHER AGENCIES

MARYLAND DNR AGENCIES:

- Wildlife & Heritage – Identify and develop restoration projects, report and map potential Ecological Significant Areas (ESA) as found during fieldwork, release programs for game and non-game species. Mapping will be done with Global Positioning Systems (GPS). Participates on the Inter-Disciplinary Team (ID Team) and assists in the development of a forest monitoring program.
- Natural Resource Police – Enforcement of natural resource laws on the forest.
- Public Lands Policy & Planning – Provides assistance in the development of plans, facilitates meetings with various management groups, develops Geographic Information System (GIS) maps for public review, and conducts deed research and boundary recovery. Also participates on the ID Team.
- Maryland Conservation Corps (MCC) – Assists in painting boundary lines, installing gates and trash removal.
- State Forest & Park Service – Participates on the ID Team.
- Chesapeake & Coastal Watershed Service – Develops watershed improvement projects, assists in the development of a forest monitoring programs and participates on the ID Team.

OTHER AGENCIES:

- DNR Contract Manager – Assists the Forest Manager in the designs and implementation of management activities on the donated portion of the forest. Also participates on the ID Team.
- Third party forest certification via annual audits
 - Sustainable Forestry Initiative (SFI)
 - Forest Stewardship Council (FSC)
- The Chesapeake Bay Foundation – Identifies sites for future water quality improvement projects and assists in the implementation by providing volunteers for reforestation.
- National Wild Turkey Federation – Establishes and maintains handicap-hunting opportunities within the forest and provides funding for habitat protection and restoration.
- US Fish & Wildlife Service – Assists in prescribed burns for Delmarva Fox Squirrel (DFS) habitat. Also assists in maintaining open forest road conditions as fire breaks.
- Maryland Forest Association - Master Loggers Program provides training in Advanced Best Management Practices for Forest Product Operators (i.e. Foresters & Loggers) workshops on the forest.
- Network with Universities and Colleges
 - Maryland Environmental Lab, Horn Point – Conducts water quality monitoring on a first order stream not influenced by agriculture. These samples will serve as a local base line for other samples taken on other Delmarva streams.
 - Allegany College – Conduct annual field tour for forestry school student’s showcasing Sustainable Forest Management practices on the forest under dual third party certification.

C. MAINTENANCE PROJECTS

Forest roads will undergo general maintenance to maintain access for forest management activities (i.e. logging, prescribed burning, and wildfire control). Interior roads within each complex will be brush hogged where possible by the MFS & the WHS. Many of the roads have grown shut and require special heavy equipment to remove the larger trees. Brushing of these roads will improve access for the public and help maintain firebreaks for communities at risk from wildfire. Recreational trails will be mowed and cleared to meet the requirements of the specific user group(s).

Forest boundary lines will be maintained using the DNR yellow band markings. Signs will be placed along the boundary lines designating the type of public access to the property. New acquisitions will be converted from their previous ownership markings to the DNR yellow band markings.

Illegal trash dumps will continue to be removed off the forest as they are discovered. The average amount of trash removed from the forest each year has been 36 tons. In our efforts to control and eradicate this issue, we will continue to coordinate with Natural Resources Police (NRP), local sheriff departments, the State Highway Administration, and County Roads departments.

D. RECREATION PROJECTS

- Host the annual Chesapeake Forest lottery for vacant tracts designated for hunt club access only. Vacant tracts are those that existing clubs opted not to continue to lease or land that has recently become available due to acquisition or right-of-ways being opened.
- Continue to explore additional Resource Based Recreational (RBR) opportunities on the forest. This may include hunting, horseback riding; water trails, hiking trails, bird watching opportunities, geocaching, etc.

- Continue work on active Recreational Trails Grants
 - Chesapeake Forest – D03 – Little Blackwater Soft Launch
 - Chesapeake Forest – D26 – Lewis/Island Pond Soft Launch
 - Pocomoke State Forest – P06 – Hudson/Tarr Handicapped Hunting Trail
 - Chesapeake Forest – W02 – Aughty Naughty Handicapped Hunting Trail
- Perform general maintenance on the existing trail system

Submit and execute Recreational Trails Grants. Appendix A contains copies of the following grant applications for Calendar Year 2015-16:

- E. Mace Smith trail marking
- Greenhill trail marking
- Furnace Town Loops

E. SPECIAL PROJECTS

- Maintain dual forest certification from the Forest Stewardship Council (FSC) and the Sustainable Forest Initiative (SFI).
- Conduct information and educational opportunities on the forest.
- Update and maintain forest information in a GIS database, which will result in a new updated forest wide field map.
- Continue the effort to inventory and protect historic sites (i.e. cemeteries, old home sites, Native American Indian sites) using GPS and GIS technology.
- Collect native genotype pond pine (*Pinus serotina*) and short-leaf pine (*Pinus echinata*) on the forest in an effort to aid future management objectives on the Pocomoke and Chesapeake Forests.
- Provide assistance to the State Tree Nursery with maintenance of Seed Orchards on the Pocomoke State Forest.

F. WATERSHED IMPROVEMENT PROJECTS

None proposed for this work plan.

G. SPECIAL WILDLIFE HABITAT PROJECTS

[CF-17-P-1]

Early-successional habitat management at Foster Tract

The existing areas along the main access road and surrounding the pond are currently comprised of predominately herbaceous vegetation. Hardwood and pine trees were planted a few years ago in some areas, but survival appears to be exceptionally low, likely due to the dry, sandy soils. Other trees species have volunteered, mainly loblolly and sweetgum, to varying extents throughout the site. These volunteer species are generally not favored by wildlife and are quite common in the surrounding forests. Therefore the WHS feels that managing these areas as early-successional habitat, in the form of herbaceous openings or pine/oak "savannas", would provide much greater benefit to wildlife than allowing them to revert to loblolly/sweetgum forests over time. This vegetation

community is not common in MD due to fire exclusion and other factors, but it provides excellent habitat for both game and non-game species.

We recommend using a combination of mechanical methods and, ideally, prescribed fire to control the woody species encroachment. Initially, many of the young loblollies and sweetgums could be removed with a skid-steer with forestry mulcher and tree shear attachments. Any desirable trees, such as oaks, hickories, or other mast-producing species, would be preserved. We estimate that this initial work could be completed in several days with the WHS's machine and labor.

Future maintenance of the early-successional habitat in these areas could be accomplished via periodic mowing or prescribed fire with WHS's equipment and labor. Prescribed fire would be preferred and provide the greatest benefit. Although the site has not been surveyed intensively, it is likely that some fire-dependent plants either currently occur or would be released on these sandy soils. Most areas have an existing trail that could easily be converted into a firebreak. However mowing every 2-3 years could also be used to control woody species if fire wasn't feasible.

H. ECOSYSTEM RESORATION PROJECTS

None proposed for this work plan.

I. MONITORING PROJECTS

The Continuous Forest Inventory (CFI) for Chesapeake Forest and Pocomoke State Forest was started in the summer of (calendar year) 2014. The CFI concluded in the winter of 2016. A summary of the results is forthcoming.